ANESTHESIA QUESTIONNAIRE - child
This questionnaire serves to brief us about your child health state. For best results of procedure please take your time to fulfill it thoroughly.

child's name		date of birth:		
weight:kg	height:cm	health insurance comp.:		
your e-mail		your phone:		
of any serious disea disease, mental and ( blood clotting disc	se, like of heart, vessel d behavioral disorder, m orders, easy bruising, pr	sp. any head injury, or has or had any signs of or respiratory disease, vascular, renal, liver of nuscular, endocrine, metabolic or hematological colonged bleeding), or any other disease? (processes)	or blocic dise	od ase
<u> </u>		cian (like neurologist, allergologist or pulmo if so, please, bring a medical report) Yo	•	
Is your child taking	any medicine on a regu	ular way? ( please bring your child's drugs v	vith vo	)II)
•	•	Y	•	No o
			es o	No o
Any previous anesthesia? Year?				No o
			∕es ∘	No (
•	•		∠es ∘	No (
	gs, food, pets, ectopic ed s are to soy, peanuts a	czema)	es o	No (
during 6 hours bef water. In the cours without solid parts Breastfeeding is all	fore procedure and no se of 6 to 2 hours befor s. Tea or water could b lowed 4 hour before p takes and thicker juice	or her, to come on an empty stomach – no drink during 2 hours before procedure, n re procedure child could drink plain water be with added sugar, without pulp or parti rocedure at last. Any other fluid, especiall es with particles or pulp could be drunk 6	neithei r, or to icles. ly fori	r ea mulas,
INFORMED CON	SENT WITH ANALO	GOSEDATION / GENERAL ANESTHES	SIA	
analgosedation. I h	nave read Parent's info	nand for her/him performing of dental properties and recommendations, I have useful ask for more information.		
Date:	Your	name and signature:		

## PARENT'S INFORMATIONS AND RECOMMENDATIONS

- For child's safety it is necessary to **come on an empty stomach no food 6 hours before procedure** and no drink **2 hours** before procedure, neither water.
- Within 6 to 2 hours before procedure child could drink plain water, or tea. Tea or water could be with added sugar, but without any solid parts.
- Breastfeeding is allowed 4 hour before procedure at the latest.
- Any other fluid, especially formulas, cow milk, milk drinks and thicker juices with particles or pulp could be drunk 6 hour before procedure at last. Not later.
- 2h before procedure your child is not allowed to drink anything.
- For more pleasant administration of oral premedication please bring your child's favorite clear drink (juice, tea, cola). Milky drinks are not convenient, nor ones with particles or pulp.
- If your child takes any drugs on a regular way, in last year suffered with any serious illness, or is in care of a specialist (neurologist, allergologist, cardiologist etc.), please bring your child's physician report. If your child takes any drugs, bring them with you.
- We appreciate if you bring light blanket for your child and his/her favorite clear non-diary drink without particles juice, tea or cola.
- Please come 30 minutes in advance at least. Time of your scheduled procedure could be changed. There is a children's corner with toys for your convenience.

## PROCEDURE OF CHILD'S ANALGOSEDATION

Procedure is intended for otherwise healthy children, in need for dental treatment, refusing to be treated without additional medication. 20-30 min before procedure sedative medication is administered in child's favorite drink. If your child's dental treatment goes only in so called **premedication**, child receives also an oral analgesics and stays partial awake during the treatment. If premedication does not insure comfortable enough condition both for your child and dental physician, another option how to safely perform dental treatment is to secure venous access, and administer **analgosedation** by this way. This means loss of consciousness and perception of pain, in a safe and controlled way, maintaining spontaneous ventilation and airway protections. This is convenient also for small preschool children. Child's vital functions are monitored during and after the procedure. Awakening to verbal contact comes in 30-120 min after procedure. Approximately 1-2h, sometimes up to 3h after procedure you could leave to home. **Full recovery is individual and it is necessary to watch your child thoroughly till next morning. Neither premedication, nor analgosedation** could be performed **in case of respiratory infection**. Please feel free to ask for more information if you would like to.

MUDr. Hynek Sechovský, mail hynek.sechovsky@gmail.com, or phone: +420 604 188 477